

The HST Key Project on the Extragalactic Distance Scale.

XXII. The Discovery of Cepheids in NGC 1326-A¹

Charles F. Prosser^{2,3}, Robert C. Kennicutt, Jr.⁴, Fabio Bresolin⁵, Abhijit Saha², Shoko Sakai²,
 Wendy L. Freedman⁶, Jeremy R. Mould⁷, Laura Ferrarese^{8,9}, Holland C. Ford¹⁰, Brad K.
 Gibson¹¹, John A. Graham¹², John G. Hoessel¹³, John P. Huchra¹⁴, Shaun M. Hughes¹⁵, Garth
 D. Illingworth¹⁶, Daniel D. Kelson¹², Lucas Macri¹⁴, Barry F. Madore^{6,17}, Nancy A. Silbermann¹⁸
 Peter B. Stetson¹⁹

¹Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by AURA, Inc. under NASA Contract No. NAS5-26555.

²NOAO, P.O. Box 26732, Tucson, AZ 85726-6732 USA

³Deceased, 15 August 1998. Charles joined the Key Project team after earning his doctorate at the University of California at Santa Cruz, and serving as a postdoctoral fellow at the Harvard-Smithsonian Center for Astrophysics. At the time of his death he had nearly completed the first draft of this paper.

⁴Steward Observatory, The University of Arizona, Tucson, AZ 85721 USA

⁵European Southern Observatory, Karl-Schwarzschild-Str. 2, Garching, Germany

⁶Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101 USA

⁷Research School of Astronomy & Astrophysics, Mount Stromlo Observatory, Weston, ACT, Australia 2611

⁸Department of Astronomy, MS 105-24, California Institute of Technology, Pasadena, CA 91125 USA

⁹Hubble Fellow

¹⁰Department of Physics & Astronomy, Johns Hopkins University, Baltimore, MD 21218 USA

¹¹CASA, University of Colorado, Boulder, CO 80309-7178 USA

¹²Department of Terrestrial Magnetism, Carnegie Institution of Washington, 5241 Broad Branch Road, NW, Washington, DC 20015, USA

¹³Department of Astronomy, University of Wisconsin, Madison, WI 53706 USA

¹⁴Harvard Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138 USA

¹⁵Institute of Astronomy, University of Cambridge, Cambridge CB3 0HA, United Kingdom

¹⁶Lick Observatory, University of California, Santa Cruz, CA 95064 USA

¹⁷NASA/IPAC Extragalactic Database and California Institute of Technology, Pasadena, CA 91125 USA

¹⁸IPAC, California Institute of Technology, Pasadena, CA 91125 USA

¹⁹Dominion Astrophysical Observatory, HIA/NRC of Canada, 5083 West Saanich Road, Victoria, BC, V8X 4M6, Canada

ABSTRACT

We report on the detection of Cepheids and the first distance measurement to the spiral galaxy NGC 1326-A, a member of the Fornax cluster of galaxies. We have employed data obtained with the Wide Field and Planetary Camera 2 on board the Hubble Space Telescope. Over a 49 day interval, a total of twelve V -band (F555W) and eight I -band (F814W) epochs of observation were obtained. Two photometric reduction packages, ALLFRAME and DoPHOT, have been employed to obtain photometry measures from the three Wide Field CCDs. Variability analysis yields a total of 17 Cepheids in common with both photometry datasets, with periods ranging between 10 and 50 days. Of these 14 Cepheids with high-quality lightcurves are used to fit the V and I period-luminosity relations and derive apparent distance moduli, assuming a Large Magellanic Cloud distance modulus $\mu_{\text{LMC}} = 18.50 \pm 0.10$ mag and color excess $E(B-V) = 0.10$ mag. Assuming $A(V)/E(V-I) = 2.45$, the DoPHOT data yield a true distance modulus to NGC 1326-A of $\mu_o = 31.36 \pm 0.17$ (random) ± 0.13 (systematic) mag, corresponding to a distance of 18.7 ± 1.5 (random) ± 1.2 (systematic) Mpc. The derived distance to NGC 1326-A is in good agreement with the distance derived previously to NGC 1365, another spiral galaxy member of the Fornax cluster. However the distances to both galaxies are significantly lower than to NGC 1425, a third Cepheid calibrator in the outer parts of the cluster.

Subject headings: galaxies: individual (NGC 1326-A) – galaxies: distances – stars: Cepheids

1. INTRODUCTION

This paper is one in a series of the *Hubble Space Telescope* Key Project on the Extragalactic Distance Scale; a recent overview of this project may be found in Kennicutt, Freedman, & Mould (1995). The primary goal is to obtain a measure for the Hubble constant (H_0) within an accuracy of $\pm 10\%$, by employing Cepheid-derived distances for nearer galaxies (redshifts generally within $\sim 1500 \text{ km s}^{-1}$) to calibrate several secondary distance indicators (Tully-Fisher relation, fundamental plane, surface brightness fluctuation method, supernovae distance indicators). In this paper we present the results from a Cepheid monitoring program of NGC 1326-A, a spiral galaxy belonging to the Fornax cluster of galaxies.

NGC 1326-A ($\alpha_{1950} = 3^{\text{h}}23^{\text{m}}$, $\delta_{1950} = -36^\circ 31'$) is the third galaxy in the Fornax cluster region which has been studied in this series. Observations of NGC 1365 and NGC 1425 have been reported by Madore et al. (1998, 1999), Silbermann et al. (1999) and Mould et al. (1999), respectively. A survey of the Fornax cluster has been reported by Ferguson (1989). Fornax is a relatively compact cluster on the sky (core radius ~ 0.7 deg), containing 350 identified members, of which most are elliptical galaxies. The heliocentric radial velocity of the Fornax Cluster is 1441 km s^{-1} , with a dispersion of 342 km s^{-1} (Madore et al. 1999). NGC 1326-A lies at a projected separation of 2.0 deg from the cluster center, with a measured heliocentric velocity of 1836 km s^{-1} (da Costa et al. 1991). The galaxy is classified as type SBm pec by de Vaucouleurs et al. (1991). It lies ~ 2.5 arcmin away from another galaxy (NGC 1326-B), and may be interacting with that system, although their radial velocities differ by over 800 km s^{-1} , which may alternatively indicate that they are a line-of-sight pair.

This paper is organized as follows. A description of the observations and data reduction is provided in §2, a description of the photometric reductions is given in §3, and the identification of variable stars given in §4. In §5, the Cepheid sample for NGC 1326-A is reviewed, and in §6 the resulting period-luminosity relations and distance modulus are presented, with further discussion and conclusions presented in §7. A series of appendices provide additional photometry, light curves, and a summary of other variable stars found.

2. OBSERVATIONS AND DATA REDUCTION

NGC 1326-A was observed using the Hubble Space Telescope Wide Field and Planetary Camera 2 (HST/WFPC2; Biretta et al. 1996) over a 49 day interval during 1997 August/September. In Figure 1, we show the WFPC2 field of view of the current observations superimposed on a V-band CCD image of NGC 1326-A, obtained with the 2.5-m telescope at the Las Campanas Observatories.

In Table 1 we list the WFPC2 observations for NGC 1326-A, giving the image identification code, UT Date of observation, the corresponding heliocentric Julian Date (mid-exposure), filter,

number of individual exposures in the epoch, and combined total exposure time. Each epoch consisted of 3 or 4 exposures taken over two orbits, to provide for better cosmic ray discrimination. Twelve epochs in *V* (F555W) and eight epochs in *I* (F814W) were obtained. As in earlier studies, the epochs roughly follow a power-law sampling distribution to reduce aliasing effects and maximize phase coverage over the ~ 10 –60 day Cepheid period range (Freedman et al. 1994, Madore & Freedman 1991). All observations were preprocessed through the standard reduction pipeline at STScI (Holtzmann et al. 1995a), which includes bias level subtraction, superbias and dark frames subtraction, and flatfielding, using the most recent calibration files available. Our own post-pipeline processing included the masking of bad columns/pixels, masking of the vignetted edges of the CCDs, and multiplication by a pixel area correction map to correct for geometric distortion in WFPC2 images (Hill et al. 1998). Finally, the calibrated images were converted to integer format for use in our photometric reduction software, after being multiplied by a factor of four in order to preserve data precision.

3. PHOTOMETRIC REDUCTIONS

As in previous studies in this series, photometric analysis of the WFPC2 images was performed independently using the software routines DAOPHOT II/ALLFRAME (Stetson 1994), and a variant of DoPHOT (Schechter et al. 1993), as described in Saha et al. (1994). In this section, we briefly summarize the separate reductions and then compare measures using a selected list of ‘secondary standard stars’ from the WFPC2 fields.

Preliminary reductions revealed very few stars on the PC chip with reliable photometry, and a full DoPHOT reduction did not yield any usable Cepheid candidates in the PC dataset. As a result, full photometric reductions in both software routines and subsequent analysis for variables were only carried out for the Wide Field CCDs WF2-WF4.

3.1. ALLFRAME Photometry

For the ALLFRAME reduction, no cosmic-ray corrections were applied to the individual galaxy frames. A master star list was created by first making two median-filtered images, one from the 40 cosmic ray split F555W images and another from the 24 F814W images. These were later summed to generate one clean frame. We adopted this method so as to include both extremely red and blue stars in the master star list. The star list was created with the automated star-finding routines from the DAOPHOT and ALLSTAR packages. This list was fed into ALLFRAME and used to extract profile-fitting stellar photometry from the 64 individual frames.

The procedures used to convert F555W and F814W instrumental magnitudes to the calibrated Landolt (1992) system are described in detail in Hill et al. (1998). For completeness, however, a brief summary follows here. The instrumental magnitudes were first brought to the $0''.5$ aperture

system of Holtzman et al. (1995). This was done by identifying 30 to 40 isolated bright stars on each chip in the median-filtered master image, and performing concentric aperture photometry on each star, using 12 apertures ranging in radius from $0''.15$ to $0''.50$. All other stars were subtracted from the original image, leaving a clean image adequate for good sky estimates. Using a growth-curve analysis provided by DAOGROW (Stetson 1990), the $0''.5$ aperture magnitude for each star was finally determined, and then compared with the corresponding PSF magnitude to calculate the aperture correction. For each filter and chip combination, the aperture correction was determined by taking the weighted mean of the aperture corrections of all the selected isolated stars. The typical star-to-star variation in aperture corrections determined for single stars is ± 0.05 mag rms, so the mean corrections are determined to an internal precision of roughly 0.01–0.02 mag, as listed in Table 2.

The transformation equation to convert the $0''.5$ aperture magnitudes, m , to the Hill et al. (1998) system, M , is expressed as:

$$M = m + 2.5 \log t + C1 + C2 * (V - I) + C3 * (V - I)^2, \quad (1)$$

where t is the exposure time and $C1$, $C2$, and $C3$ are constants. The coefficients $C2$ and $C3$ for the color-corrections were adopted from Holtzman et al. (1995); they are the same for all four chips. The constant $C1$ consists of several terms: the long-exposure WFPC2 magnitude zero points, the ALLFRAME magnitude zero point, a correction for multiplying the original image by four before converting them to integers, a gain ratio term due to the difference between the gain settings used for NGC 1326-A and for the Holtzman et al. data (7 and 14 respectively), and a correction for the pixel area map which was normalized differently from Holtzman et al. (1995). Each term is discussed in detail by Hill et al. (1998). Values of the $C1$ terms are listed in Table 2. As was the case for other papers in this series, we have adopted the “long exposure” zeropoints of Hill et al. (1998) for both the ALLFRAME and DoPHOT photometry.

Recently Stetson (1998) has reanalyzed the zeropoint calibration of WFPC2 taking into account possible second-order charge transfer effects. In order to maintain consistency with previous papers in this series we have not adopted the corrections derived in Stetson (1998), adopting instead the Hill et al. (1998) zeropoints with the 0.05 mag correction for charge transfer effects. Any differences between the two scales should be very small in comparison to the other uncertainties in the Cepheid distance to NGC 1326-A. A new analysis of the photometric zeropoint is currently being carried out by P. Stetson, and will be reported in a future paper.

3.2. DoPHOT Photometry

The DoPHOT reduction followed the procedures described in Ferrarese et al. (1996, 1998). Additional detailed discussion of the procedure may also be found in Saha et al. (1996). We

briefly summarize the reduction procedure here, and mention aspects specific to the current study of NGC 1326-A.

Prior to running DoPHOT, the 3-4 individual exposures at each epoch in Table 1 were coadded into a single image, with cosmic ray events identified and rejected in the process. After coalignment of the images from each epoch, master F555W and F814W frames were derived from median filtering of the combined epoch observations. DoPHOT was then run on these deep images to produce a master object list, which was then transformed and applied as input when running DoPHOT on each separate epoch. The output PSF photometry (m_{psf}) from DoPHOT is then scaled by the exposure time (t) and calibrated onto the Holtzmann et al. (1995) $0''.5$ ‘ground system’ (m_{grd}) through the use of aperture correction (AC) and zero point (ZP) terms:

$$m_{\text{grd}} = m_{\text{psf}} + 30.0 + 2.5\log(t) + \text{AC} + \text{ZP} \quad (2)$$

A different approach to calculating the aperture corrections is used in the DoPHOT reductions (Saha et al. 1996). The positional variation in the corrections is determined from external data with a clean background. Due to the paucity of bright isolated stars in NGC 1326-A, the mean aperture corrections were derived from single-epoch observations of the Local group dwarf spheroidal galaxy Leo I, as described in Saha et al. (1996). This procedure has been tested extensively for other galaxies in the project and has been shown to reproduce the aperture corrections within the uncertainties over a wide range of WFPC2 data. The coefficients used to derive these AC values are given in Table 3, along with the zero point (ZP) values used. The ZP values include corrections for the gain ratio term among the different chips. The rms scatter in AC measurements for single stars, after fitting of the spatial variation, is ± 0.035 mag rms, and the rms uncertainties in the average corrections are roughly an order of magnitude smaller.

The mean F555W and F814W magnitudes of the Cepheids and the secondary standard stars were converted to standard V and I magnitudes, using the color correction coefficients from Holtzmann et al. (1995, Table 3):

$$V = \text{F555W} - 0.045(\text{F555W} - \text{F814W}) + 0.027(\text{F555W} - \text{F814W})^2, \quad (3)$$

$$I = \text{F555W} - 0.067(\text{F555W} - \text{F814W}) + 0.025(\text{F555W} - \text{F814W})^2. \quad (4)$$

3.3. Comparison of DoPHOT and ALLFRAME Photometry

To compare the DoPHOT and ALLFRAME magnitudes, we selected stars in each the 3 WF chips which were relatively isolated, showed low formal errors in their magnitude measures, were distributed over the CCD, and covered a magnitude range in common with the detected Cepheid variables. The locations and DoPHOT photometry of these ‘secondary standard’ stars are given in Appendix A. Figure 2 shows plots of the V and I magnitude differences (DoPHOT

– ALLFRAME) for the WF2-WF4 chips. The corresponding mean differences and errors are given in Table 4. The magnitudes of the secondary standard stars show reasonable consistency, especially when taking into account the relatively large distance and high crowding level in this galaxy. The Cepheids themselves show somewhat larger differences, reflecting the fact that the Fornax cluster lies near the limit at which accurate crowded-field photometry can be obtained with WFPC2.

4. IDENTIFICATION OF VARIABLE STARS

In the case of the ALLFRAME data set the search for variables was carried out with a variation of the correlated variability test of Welch & Stetson (1993), as implemented in the algorithm for the automated detection of Cepheids described by Stetson (1996). This method, which relies on the temporal correlation of magnitude residuals of real variables, takes advantage of the CR split nature of the HST observations.

Variability analysis of the DoPHOT data set was performed on the F555W observations, following the general technique detailed in Saha & Hoessel (1990). In order to be flagged as a variable, a star had to be measured in at least 8 out of the 12 V -band epochs. Stars with nearby neighbors (within 2 pixels) that contributed more than 50% of the total light were rejected. Variable stars were identified using a simple reduced χ^2 criterion, with the requirement that the variability be significant at the >99% confidence level, or $\chi^2 > 8$ (Ferrarese et al. 1998). Stars with confidence levels above 99% but with lower χ^2 values were tested further for periodicity, using a variant of the Lafler-Kinman (1965) method of phase dispersion minimization. Stars with $\Lambda \geq 2.8$ were flagged as candidate variables, where Λ is as defined in Lafler & Kinman (1965), and implemented in Saha & Hoessel (1990).

NGC 1326-A is one of the most distant galaxies in the Key Project, and this combined with its relatively low luminosity yields a smaller potential sample of candidate variables than most of the other galaxies in the project. For this study, we performed an additional check on the significance of the variability, by performing random redistribution of the observational dataset to check directly the random probability of detecting each candidate as a variable. We performed 100 random resamplings for each candidate, and applied the reduced χ^2 and Λ tests described above to these data. This ‘bootstrapping’ technique allowed us to eliminate numerous spurious variables which passed the variability tests due to a single outlying observation, usually from a bad pixel or cosmic ray event. A final check for spurious variables was made by visually inspecting and blinking the images for each candidate.

Periods for the candidate variable stars were determined by phasing the data for all periods between 3 and 100 days, and applying the Lafler-Kinman phase dispersion minimization criterion. Because of the logarithmic sampling of the data, period aliases are not a serious problem, and in the rare ambiguous cases the data were checked for spurious observations and the variability

analysis was performed interactively. Serious aliasing was often a problem for stars with periods longer than our observing window (49 days), so only those Cepheids with $P < 50$ days are analyzed in this paper.

5. CEPHEID VARIABLE STARS

5.1. Cepheid Sample

The final list of Cepheid variable stars in NGC 1326-A is composed of stars which meet the variability criteria in both of the ALLFRAME and DoPHOT data sets. When matching the candidate lists we also performed several other checks on the properties of the stars. Stars with lightcurve shapes that clearly differ from Cepheids, stars with low variability amplitude or flat-bottomed lightcurves, those with highly abnormal colors, or those with strong variability in V but little or none in I were all rejected from the final list. These tests were aimed at rejecting non-Cepheid variable stars and Cepheids with unresolved companions. Stars located in crowded regions were also rejected, according to the criteria given in §4. None of these rejected candidates were included in the final Cepheid sample, but a list of indeterminate variables and long-period variables is given in Appendix B.

Adopting these criteria yielded 17 high-quality Cepheids in common between the DoPHOT and ALLFRAME lists. These stars, along with their astrometric positions are listed in Table 5, in order of decreasing period. The absolute positions are accurate to within the pointing accuracy of HST ($\sim 1''$), while the relative positions should be accurate to approximately $0.''2$. Finder charts for the Cepheids (as well as the other variable stars) are given in Figure 3, and enlarged images of the regions surrounding each Cepheid are given in Figure 4. DoPHOT and ALLFRAME mean photometry is given in Tables 6 and 7, respectively. These tables include two candidates, C8 and C10, which yielded high-quality lightcurves with DoPHOT only. Such inconsistency between the two programs is usually caused by semi-resolved companions to the Cepheids, so as a precaution these are listed but are not included in the fitting of the period-luminosity relation. Individual DoPHOT magnitudes in F555W and F814W (and ALLFRAME V and I magnitudes) at each epoch are provided in Appendix C¹. Finally, the phased lightcurves of the Cepheids in F555W and F814W are shown in Figure 5.

¹Additionally, these values may be obtained through the HST Key Project Archives at <http://www.ipac.caltech.edu/H0kp>.

5.2. Mean Magnitudes

Tables 6 and 7 list both intensity averaged (‘av’ subscript) and phase weighted (‘ph’ subscript) mean magnitudes for each Cepheid, defined as

$$m_{\text{av}} = -2.5 \log_{10} \sum_{i=1}^n \frac{1}{n} 10^{-0.4m_i}, \quad (5)$$

$$m_{\text{ph}} = -2.5 \log_{10} \sum_{i=1}^n 0.5(\phi_{i+1} - \phi_{i-1}) 10^{-0.4m_i}, \quad (6)$$

where n is the total number of observations and m_i and ϕ_i are the magnitude and phase of the i th observation in order of increasing phase. The variability levels of reduced χ^2 for DoPHOT and σ for ALLFRAME are provided in the last columns of the tables. In the case of the F814W (I) observations, due to the fewer phase points available the phase weighted means have been adjusted by the small correction factor based on the V to I amplitude ratio, following the procedure described in Freedman et al. (1994). As may be expected, differences between m_{av} and m_{ph} are larger when the phase coverage is less uniform.

The periods determined from the DoPHOT and ALLFRAME fitting show very good agreement, usually within 5% or better. The mean uncertainty in the intensity and phase-weighted mean magnitudes is ± 0.1 mag, which is comparable to the consistency of the DoPHOT and ALLFRAME magnitudes, as summarized in Table 4. This level of agreement in magnitude scales is lower than in most of the other galaxies in the Key Project sample, but is comparable to the errors found previously in galaxies near the distance limit of this project, e.g., NGC 4414 (Turner et al. 1998), NGC 1365 (Silbermann et al. 1999), and NGC 4548 (Graham et al. 1999). We have included these uncertainties in the total error budget for the distance to NGC 1326-A (§6).

Figure 6 shows the V vs. $V - I$ color magnitude diagram for NGC 1326-A, based on the DoPHOT photometry of the three WF chips. Phase-averaged mean magnitudes for the Cepheids are overplotted, along with the expected $P = 10 - 60$ day locus line for zero reddening and an assumed distance modulus of 31.4 mag.

6. PL RELATIONS AND THE DISTANCE TO NGC 1326-A

6.1. PL Relations and Apparent Distance Moduli

Apparent distance moduli to NGC 1326-A in V and I were determined using the final set of Cepheids listed in Tables 5–7, and the calibrating PL relations from Madore & Freedman (1991):

$$M_V = -2.76(\log P - 1.0) - 4.16 \quad (7)$$

$$M_I = -3.06(\log P - 1.0) - 4.87 \quad (8)$$

These calibrations are based on a sample of 32 Cepheids in the LMC, and assume an LMC true modulus and average line-of-sight reddening of 18.50 ± 0.10 and $E(B - V) = 0.10$ mag respectively. As in all other papers in this series, the slopes of the PL relations are fixed to the values given in equations (5) and (6), and only the mean shift from the LMC relation is fitted. This minimizes any biases which might be introduced by incompleteness at short periods in the NGC 1326-A sample.

The PL relations for the NGC 1326-A Cepheids from the DoPHOT data are shown in Figure 7. The relations were fitted using 14 of the Cepheids in Tables 5–7, shown as solid points in Figure 7. Stars C08 and C10 were not used in the fit because they were identified cleanly only in the DoPHOT data set. C11 and C19 were not fitted because they show very anomalous colors in the DoPHOT data, differing by more than half a magnitude from the expected colors for Cepheids of their respective periods, compared to an rms photometric error of ~ 0.15 mag in $V - I$ (C19 also has a period below 10 days). Such anomalous colors are often a sign of an unresolved blend with a companion star, so as a conservative measure C11 and C19 were excluded from the PL fitting. C04 was also rejected because it lies more than 3σ above the mean PL relation, probably due to an unresolved blend with another star, which would also account for its low amplitude. The Cepheids which were not included in the fits are shown as open circles in Figure 7. The 14 stars used to fit the PL relation have periods in the range 10–50 days.

The fits to the V and I PL relations were performed as described in Freedman et al. (1994a). Fitting the DoPHOT periods and phase weighted magnitudes yields apparent moduli $\mu_V = 31.36 \pm 0.06$ mag, $\mu_I = 31.36 \pm 0.06$ mag, and $E(V - I) = 0.00 \pm 0.05$ mag, where the quoted uncertainties refer to the rms dispersion in the mean PL fits only. The corresponding moduli from the ALLFRAME data set are $\mu_V = 31.39 \pm 0.07$ mag, $\mu_I = 31.34 \pm 0.06$ mag, and $E(V - I) = 0.05 \pm 0.03$ mag. The best fitting PL relations are shown by the solid lines in Figure 7. The dashed lines, drawn at ± 0.54 mag for the V PL plot, and ± 0.36 mag for the I PL plot, represent the 2σ scatter of the best fitting PL relation for the LMC Cepheids (Madore & Freedman 1991).

6.2. Extinction and True Distance Modulus

The true distance modulus to NGC 1326-A, μ_0 , is given by

$$\mu_0 = \mu_V - A(V) = \mu_I - A(I) \quad (9)$$

where the V and I band absorption coefficients $A(V)$ and $A(I)$ obey the relation $A(V)/E(V - I) = 2.45$. This is consistent with the extinction laws Dean, Warren, & Cousins (1978),

Cardelli, Clayton & Mathis (1989), and Stanek (1996) and assumes $R_V = A(V)/E(B - V) = 3.3$. Applying this to the DoPHOT PL fits given above yields a true distance modulus $\mu_0 = 31.36 \pm 0.11$ mag, again with the quoted uncertainty only reflecting the formal dispersion in the individual Cepheid distance moduli. The ALLFRAME data give a similar value for the true modulus, $\mu_0 = 31.27 \pm 0.08$ mag. The corresponding distances for the ALLFRAME and DoPHOT fits are 18.7 Mpc and 17.9 Mpc, respectively. The uncertainties associated with these distance are discussed below.

6.3. Error Budget

The error budget in the determination of the true distance modulus is summarized in Table 8. To maintain consistency with the previous papers in this series we closely follow the analysis of Phelps et al. (1998). This paper contains a detailed discussion and justification for the individual error terms, and here we only summarize those aspects which are specific to NGC 1326-A.

The first error terms in Table 8 (*a-c*, *S1*) are those arising from the LMC-based calibration of the PL relation, including errors in the assumed LMC distance and uncertainties in the calibrating Cepheid photometry. Because this uncertainty affects all Key Project galaxies exactly the same way, it is of a systematic nature. The second set of terms (*d-e*, *R1*) include uncertainties in the LMC photometric zeropoints, as discussed in detail by Hill et al. (1998) and Stetson et al. (1998). These introduce a significant uncertainty in the true distance modulus because of the strong propagation of color errors in the dereddened modulus. Uncertainties in the PL fits and reddening to NGC 1326-A introduce an additional random error (*f-g*, *R2*).

Another source of systematic uncertainty in the distance moduli is the uncertain dependence of the Cepheid PL relation on metal abundance (Kennicutt et al. 1998). However this is not a serious problem for NGC 1326-A. Spectrophotometry of five HII regions in the galaxy by Kennicutt et al. (1999) yields a mean oxygen abundance $12 + \log(O/H) = 8.48 \pm 0.16$ when calibrated on the scale of Zaritsky, Kennicutt, & Huchra (1994). This is virtually identical to the mean value for the LMC of 8.50 ± 0.1 when calibrated on the same system. Consequently any metallicity effect should be minimal, and the error term (*S2*) in Table 8 only reflects the effect of the uncertainty in the metal abundance itself, when propagated with the Cepheid metallicity dependence in Kennicutt et al. (1998).

Combining these error terms in quadrature with the PL fits from the DoPHOT photometry yields final values for the distance modulus of NGC 1326-A of $\mu_0 = 31.36 \pm 0.17$ (random) ± 0.13 (systematic) mag. The corresponding distance is 18.7 ± 1.5 (random) ± 1.2 (systematic) Mpc. The ALLFRAME analysis yields a distance that is 3.3% lower ($\mu_0 = 31.27$, $d = 17.9$ Mpc), with the same uncertainties. In this case we preferentially quote the DoPHOT distance, because the PL relations are somewhat tighter and the fitted distance modulus is less sensitive to whether the shortest-period variables are excluded. However both sets of distances are consistent within their

quoted errors.

7. DISCUSSION

NGC 1326-A is a relatively unstudied galaxy, and no previous distance measurements could be found in the literature. An extensive literature exists on the distance of the Fornax cluster, of course, as discussed in Madore et al. (1999). The distance of 18.7 Mpc that we derive to NGC 1326-A, although uncertain at the 10% level, is in excellent agreement with the distance of 18.2 ± 1.7 (random) ± 1.5 (systematic) Mpc derived to another Fornax cluster member, NGC 1365, based on measurements of 52 Cepheids (Silbermann et al. 1999). However the distances we derive to NGC 1326-A and NGC 1365 are significantly smaller than that derived for the third Fornax cluster galaxy in our sample, NGC 1425, at 22.2 ± 1.7 (random) ± 1.8 (systematic) Mpc (Mould et al. 1999). A detailed comparison of these distances and their interpretation is given in Mould et al. (1999).

We acknowledge the assistance of Matthew Trewhella in retrieving from archive the CCD image used in Figure 1. We also thank the anonymous referee for comments which improved the paper significantly. We gratefully acknowledge the support of the NASA and STScI support staff. Support for this work was provided by NASA through grant G0-2227-87A from STScI.

A. SECONDARY STANDARDS IN THE NGC 1326-A FIELD

Tables 9–11 list photometry for the stars selected as secondary standard stars in each of the WF chips, as described in §3.3 and used in the comparison shown in Figure 2. The tables list for each star the DoPHOT (x,y) coordinates on the CCD chips, DoPHOT F555W and F814W ground system magnitudes, the corresponding calibrated V and I DoPHOT magnitudes, and the calibrated V and I ALLFRAME magnitudes.

B. OTHER VARIABLE STARS IN NGC 1326-A

Tables 12–13 list identification information and photometry for 17 miscellaneous variable stars in NGC 1326-A. These stars include Cepheids which were excluded from the final sample because of uncertain periods, large magnitude errors, or inconsistencies between the ALLFRAME and DoPHOT data sets, long-period variables with periods longer than our sampling window, or other types of variable stars, as described in §4.2. Table 12 contains positional information and notes which provide additional information. Table 13 lists DoPHOT photometry and period information (when available) for these stars.

C. INDIVIDUAL EPOCH PHOTOMETRY OF CEPHEIDS

Tables 14–15 list the DoPHOT F555W and F814W (ground system) photometry for the final sample of Cepheids on an epoch-by-epoch basis. Tables 16–17 list the corresponding ALLFRAME photometry.

REFERENCES

- Biretta, J., et al. 1996, Wide Field and Planetary Camera 2 Instrument Handbook, version 4.0 (Baltimore: STScI)
- da Costa, L.N., Pellegrini, P.S., Davis, M., Meiksin, A., Sargent, L.W., & Tonry, J.L. 1991, ApJS, 75, 935
- de Vaucouleurs, G., et al. 1991, Third Reference Catalogue of Bright Galaxies (New York: Springer-Verlag)
- Ferguson, H.C. 1989, AJ, 98, 367
- Ferrarese, L., et al. 1998, ApJ, 507, 655 (Paper XII)
- Freedman, W.L., et al. 1994, ApJ, 427, 628 (Paper I)
- Graham, J.A., et al. 1999, ApJ, in press (Paper XX)
- Hill, R., et al. 1998, ApJ, 496, 648 (Paper V)
- Holtzmann, J., et al. 1995a, PASP, 107, 156
- Holtzmann, J., et al. 1995b, PASP, 107, 1065
- Kennicutt, R.C., Freedman, W.L., & Mould, J.R. 1995, AJ, 110, 1476
- Kennicutt, R.C., et al. 1998, ApJ, 498, 181 (Paper XIII)
- Kennicutt, R.C., Huchra, J.P., Macri, L.M., & Mould, J.R. 1999, in preparation
- Lafler, J., & Kinman, T.D. 1965, ApJS, 11, 216
- Madore, B.F., & Freedman, W.L. 1991, PASP, 103, 933
- Madore, B.F., et al. 1998, Nature, 395, 47
- Madore, B.F., et al. 1999, ApJ, 515, 29 (Paper XV)
- Mould, J.R. et al. 1999, ApJ, submitted (Paper XXI)
- Phelps, R.L., et al. 1998, ApJ, 500, 763 (Paper IX)
- Saha, A., & Hoessel, J.G. 1990, AJ, 99, 97
- Saha, A., Labhardt, L., Schwengeler, H., Macchetto, F.D., Panagia, N., Sandage, A., & Tammann, G.A. 1994, ApJ, 425, 14

- Saha, A., Sandage, A., Labhardt, L., Tammann, G.A., Macchetto, F.D., & Panagia, N. 1996, ApJ 46, 55
- Schechter, P.L., Mateo, M., & Saha, A. 1993, PASP, 105, 1342
- Silbermann, N.A., et al. 1999, ApJ, 515, 1 (Paper XIV)
- Stetson, P.B. 1990, PASP, 102, 932
- Stetson, P.B. 1994, AJ, 106, 205
- Stetson, P.B. 1996, PASP, 108, 851
- Stetson, P.B. 1998, PASP, 110, 1118
- Stetson, P.B. et al. 1998, ApJ, 508, 491 (Paper XVI)
- Turner, A. et al., 1998, ApJ, 505, 207 (Paper XI)
- Welch, D.L., & Stetson, P.B. 1993, AJ, 105, 1813
- Zaritsky, D., Kennicutt, R.C., & Huchra, J.P. 1994, ApJ, 420, 87

Figure Captions

Fig. 1.— V-band ground based image of NGC 1326-A, with the approximate dimensions of the HST WFPC2 field of view overplotted. The bright galaxy on the edge of the image is NGC 1326-B. North is to the top and east to the left.

Fig. 2.— Comparison between the DoPHOT/ALLFRAME V and I magnitudes among the secondary standard stars from the three WF CCDs.

Fig. 3.— Median-filtered V-band images of the four WFPC2 chips. The locations of the Cepheids and other variable stars recovered in this study are indicated by circles, and are labeled as listed in Tables 5 and 12 . In each frame, the pyramid for the WFPC2 camera is in the lower lefthand corner.

Fig. 4.— Expanded finder charts for the Cepheids (Table 5) and the other variable stars (Table 12). Each panel is 50×50 pixels on the WFC, corresponding to $5''$ square on the sky.

Fig. 5.— Light curves for the Cepheid variables, based on the DoPHOT photometry. The V magnitudes are plotted as solid points, the I magnitudes as open squares. The errorbars reported by DoPHOT are indicated.

Fig. 6.— V vs. $V - I$ color magnitude diagram for NGC 1326-A, based on DoPHOT photometry from median filtered images of the four WFPC2 chips. The recovered Cepheid sample is shown by large points. The expected ridge line for Cepheids with $10 \leq P \leq 60$ days and zero reddening is drawn for reference, assuming an apparent distance modulus of 31.4 mag. Photometric error bars for a typical single epoch of observation are also shown.

Fig. 7.— PL relations in V (top) and I (bottom) for Cepheids in NGC 1326-A, based on the DoPHOT photometry. Cepheids included in the fitting are shown as solid points, while open circles denote Cepheids that were rejected from the fit, as described in §6.1. The solid line shows the fitted PL relation, while the dashed lines show the 2σ intrinsic width of the PL relation in the LMC.

Table 1. NGC 1326-A Journal of Observations

STScI Rootname ID	1997 UT Date	Average HJD	Filter	No. Exp.	Exposure Time (sec)
U34L2601M	AUG06	2450667.0952	F555W	3	3400
U34L2701M	AUG09	2450670.1196	F555W	3	3400
U34L2801M	AUG12	2450673.0080	F555W	3	3900
U34L2901M	AUG15	2450675.9062	F555W	4	5200
U34L3001M	AUG18	2450679.2000	F555W	4	5200
U34L3101R	AUG23	2450684.3708	F555W	3	3900
U34L3201R	AUG27	2450687.7434	F555W	4	5200
U34L3301R	SEP01	2450693.3174	F555W	3	3900
U34L3401O	SEP05	2450697.2925	F555W	4	4600
U34L3501R	SEP12	2450704.2744	F555W	3	3400
U34L3601R	SEP17	2450709.3821	F555W	3	3400
U34L3701R	SEP24	2450716.1699	F555W	3	3400
...	...	...	...	...	...
U34L2604M	AUG06	2450667.1765	F814W	3	3600
U34L2704M	AUG09	2450670.2008	F814W	3	3600
U34L2804M	AUG12	2450673.0917	F814W	3	3900
U34L3104R	AUG23	2450684.4542	F814W	3	3900
U34L3304R	SEP01	2450693.4007	F814W	3	3900
U34L3504R	SEP12	2450704.3557	F814W	3	3600
U34L3604O	SEP17	2450709.4633	F814W	3	3600
U34L3704R	SEP24	2450716.2515	F814W	3	3600

Table 2. Coefficients Used for ALLFRAME Photometry

Chip	Filter	Stars	Aperture Correction Zero point ^a	WFPC2 Coeff. ^b	C1
WF2	F555W	51	-0.027 ± 0.017	22.522	-0.984
WF2	F814W	51	$+0.006 \pm 0.006$	21.616	-1.857
WF3	F555W	22	$+0.015 \pm 0.006$	22.530	-0.934
WF3	F814W	22	$+0.023 \pm 0.011$	21.638	-1.818
WF4	F555W	14	$+0.009 \pm 0.009$	22.506	-0.964
WF4	F814W	14	-0.005 ± 0.011	21.609	-1.875

^aWFPC2 ZP values from Hill et al. (1998, Table 7).

^b ALLFRAME instrumental F555W/F814W magnitudes (m_{inst}) are obtained from the fitted magnitudes (m_{fit}) using the relation: $m_{\text{inst}} = m_{\text{fit}} + 2.5 \times \log t + C$, where $C = -25.00 + 0.016 + 2.5 \times \log 4 + AC + ZP$, and t is the exposure time in seconds. The terms in the C coefficient include: -25.00 as the ALLFRAME zeropoint, $+0.016$ due to the applied pixel area normalization, a $2.5 \log 4$ term since the data have been converted to integer format and multiplied by a factor of 4, and aperture correction (AC) and WFPC2 zeropoint (ZP) terms as given above.

Table 3. DoPhot Aperture Correction Coefficients^a and Zeropoints^b

Filter	Chip	CF0	CF1	CF2	CF3	CF4	CF5	ZP
F555W	PC1	−0.880	−6.43E−05	+1.49E−04	−2.00E−07	−3.66E−07	+5.31E−07	−7.677
F555W	WF2	−0.702	+1.67E−04	+2.63E−04	−2.34E−07	−3.02E−07	+4.66E−08	−7.586
F555W	WF3	−0.614	+1.16E−06	−1.02E−04	−1.24E−07	−4.39E−08	+2.67E−07	−7.599
F555W	WF4	−0.682	+3.37E−04	+1.34E−04	−4.68E−07	−3.46E−07	+2.15E−07	−7.601
F814W	PC1	+1.065	+4.33E−05	+1.46E−04	−2.29E−07	−3.11E−07	+3.98E−07	−8.548
F814W	WF2	−0.740	+4.31E−05	+1.07E−04	−1.71E−07	−2.32E−07	+2.18E−07	−8.416
F814W	WF3	−0.763	+4.00E−05	+4.78E−05	−3.78E−08	−6.64E−08	+5.68E−09	−8.472
F814W	WF4	−0.764	+1.92E−04	+1.27E−04	−2.63E−07	−2.52E−07	+6.88E−08	−8.471

^aAperture correction coefficients determined from observation of Leo I dwarf galaxy field.

Aperture correction value: $AC = CF0 + CF1 \times X + CF2 \times Y + CF3 \times X^2 + CF4 \times Y^2 + CF5 \times X \times Y$.

^bDoPhot zeropoints place photometry onto WFPC2 Ground System of Holtzmann et al. (1995).

Table 4. Comparison of DoPHOT and ALLFRAME Photometry

Set	Chip	N(stars)	$\Delta V \pm \sigma_V$	$\Delta I \pm \sigma_I$
Secondary Standards	WF2	55	−0.04 (0.01)	−0.02 (0.02)
	WF3	62	−0.07 (0.01)	−0.07 (0.01)
	WF4	20	+0.04 (0.03)	+0.05 (0.01)
Cepheids	WF2	10	−0.12 (0.08)	+0.01 (0.15)
	WF3	6	−0.03 (0.06)	−0.11 (0.12)
	WF4	1	+0.02 (...)	+0.27 (...)

Note: The magnitude differences listed are in the sense DoPHOT − ALLFRAME, and the quoted errors represent the standard deviation of the mean difference.

Table 5. Astrometry for NGC 1326-A Cepheids

ID ^a	PL ^b	X ^c	Y ^c	RA(J2000) ^d	DEC(J2000) ^d	Chip
C1	+	185.15	385.24	3:25:05.91	−36:21:37.4	WF2
C2	+	547.56	542.25	3:25:04.95	−36:20:59.9	WF2
C3	+	97.11	690.93	3:25:08.30	−36:21:24.3	WF2
C4		610.93	493.90	3:25:10.40	−36:22:07.5	WF3
C5	+	225.81	312.48	3:25:07.01	−36:22:18.6	WF3
C6	+	463.41	510.73	3:25:05.20	−36:21:08.3	WF2
C7	+	412.55	527.10	3:25:05.58	−36:21:11.1	WF2
C8		299.18	94.68	3:25:06.31	−36:21:57.3	WF3
C9	+	224.76	274.70	3:25:05.01	−36:21:41.6	WF2
C10		356.75	221.03	3:25:07.34	−36:22:03.2	WF3
C11		421.88	269.38	3:25:08.01	−36:22:02.7	WF3
C12	+	459.80	303.39	3:25:03.93	−36:21:22.0	WF2
C13	+	239.58	302.26	3:25:07.04	−36:22:16.9	WF3
C14	+	192.82	88.33	3:25:05.61	−36:22:03.7	WF3
C15	+	200.38	238.98	3:25:04.91	−36:21:45.8	WF2
C16	+	419.14	189.51	3:25:03.43	−36:21:32.5	WF2
C17	+	435.19	394.31	3:25:08.76	−36:22:11.3	WF3
C18	+	242.22	221.38	3:25:04.32	−36:22:37.0	WF4
C19		485.98	662.95	3:25:06.03	−36:20:56.7	WF2

^aCepheids are ordered by increasing period.

^bPL column is flagged if the cepheid is used in fit to the period-luminosity relation for NGC 1326-A.

^cX and Y coordinates are derived from UT 1997 August 06 epoch.

^dRA and DEC are derived using the IRAF task `stsdas.toolbox.imgtools.xy2rd`, STSDAS V2.0.1.

Table 6. DoPHOT Mean Photometry for NGC 1326-A Cepheids

ID	P	P _{err}	χ^2_r	log P	F555W _{av}	F814W _{av}	F555W _{ph}	F814W _{ph}	V _{ph}	I _{ph}	V–I
C01	50.	(6)	28.9	1.70	25.11	24.18	25.06	24.16	25.04	24.12	0.92
C02	40.3	(4)	18.0	1.61	25.56	24.82	25.68	24.85	25.66	24.81	0.85
C03	34.8	(3)	43.8	1.54	25.52	24.83	25.58	24.87	25.56	24.83	0.73
C04	34.8	(3)	11.9	1.54	25.11	24.21	25.06	24.14	25.04	24.10	0.94
C05	34.8	(3)	8.5	1.54	26.02	25.08	26.03	25.17	26.01	25.13	0.88
C06	32.1	(3)	52.8	1.51	25.73	24.95	25.73	24.93	25.71	24.89	0.82
C07	29.8	(3)	24.5	1.47	25.95	25.12	25.93	25.04	25.91	25.00	0.91
C08	27.8	(3)	24.1	1.44	25.88	24.97	25.84	25.02	25.82	24.98	0.84
C09	26.6	(3)	18.9	1.42	26.07	25.33	26.08	25.38	26.06	25.34	0.72
C10	21.6	(2)	12.2	1.33	26.27	25.23	26.27	25.15	26.25	25.04	1.14
C11	21.2	(2)	1.2	1.33	26.33	25.00	26.34	24.97	26.34	24.93	1.40
C12	20.9	(2)	13.4	1.32	26.43	25.87	26.44	25.93	26.42	25.90	0.52
C13	20.5	(2)	4.5	1.31	26.56	25.36	26.56	25.38	26.54	25.34	1.20
C14	19.3	(1)	10.4	1.29	26.80	25.78	26.66	25.83	26.64	25.79	0.85
C15	17.5	(1)	9.5	1.24	26.28	25.30	26.31	25.41	26.29	25.37	0.92
C16	15.4	(1)	4.1	1.19	27.07	26.21	27.06	26.08	27.04	26.04	1.00
C17	12.4	(0.7)	13.3	1.09	26.66	25.95	26.66	26.03	26.64	26.00	0.64
C18	11.4	(0.8)	4.2	1.06	26.66	26.00	26.71	26.20	26.69	26.17	0.52
C19	9.7	(0.3)	8.1	0.99	26.73	26.35	26.77	26.41	26.76	26.39	0.37

Table 7. ALLFRAME Mean Photometry for NGC 1326-A Cepheids

ID	P	P _{err}	Var. σ	log P	V _{av}	I _{av}	V _{ph}	I _{ph}	V–I _{ph}
C01	47.4	(0.7)	2.8	1.68	25.22	24.23	25.18	24.26	0.92
C02	39.5	(0.4)	2.0	1.60	25.69	24.81	25.77	24.83	0.94
C03	35.9	(0.7)	3.2	1.56	25.44	24.72	25.54	24.75	0.79
C04	33.7	(0.6)	1.8	1.53	25.19	24.23	25.14	24.20	0.94
C05	32.0	(1.2)	1.2	1.51	26.03	25.18	26.08	25.23	0.85
C06	30.8	(0.3)	2.4	1.49	25.75	24.97	25.77	24.93	0.84
C07	30.1	(0.4)	2.8	1.48	26.01	25.15	25.99	25.13	0.86
C08									
C09	25.2	(0.4)	1.4	1.40	26.17	25.31	26.18	25.33	0.85
C10									
C11	20.9	(0.4)	0.8	1.32	26.30	25.15	26.33	25.18	1.15
C12	19.7	(0.3)	1.3	1.29	26.55	25.81	26.56	25.71	0.85
C13	19.6	(0.5)	0.5	1.29	26.59	25.52	26.60	25.61	0.99
C14	20.3	(0.6)	0.9	1.31	26.63	25.75	26.58	25.76	0.82
C15	17.5	(0.3)	1.5	1.24	26.38	25.35	26.40	25.37	1.03
C16	14.7	(0.2)	0.6	1.17	27.20	26.21	27.20	26.20	1.00
C17	12.3	(0.1)	0.6	1.09	26.64	25.97	26.67	26.00	0.67
C18	10.3	(0.2)	0.3	1.01	26.69	25.83	26.67	25.89	0.78
C19	9.8	(0.1)	0.8	0.99	26.85	26.17	26.84	26.09	0.75

Table 8. Error Budget in the Distance for NGC 1326-A

Source of Uncertainty	Error (mag)
Cepheid Period Luminosity Calibration	
(a) LMC True Distance Modulus	± 0.10
(b) V-band LMC PL Zero Point	± 0.05
(c) I-band LMC PL Zero Point	± 0.03
(S1) LMC PL Systematic Uncertainty	± 0.12
HST Photometric Calibration	
(d) WFPC2 V-band Photometry Zero Point	± 0.05
(e) WFPC2 I-band Photometry Zero Point	± 0.05
(R1) HST Calibration Uncertainty	± 0.15
NGC 1326-A Modulus	
(f) NGC 1326-A V-band PL Fitting	± 0.06
(g) NGC 1326-A I-band PL Fitting	± 0.06
(R2) Distance Modulus Uncertainty	± 0.08
Metallicity Effect	
(S2) Uncertainty from Metallicity Difference (LMC vs. NGC1326-A)	± 0.05
Total Uncertainty	
(R) Random Errors (R1+R2)	± 0.17
(S) Systematic Errors (S1+S2)	± 0.13

Note: S1 is derived by combining (a), (b), and (c) in quadrature. R1 is derived by combining (d) and (e) in quadrature, but weighted by the reddening coefficients, $R1 = [(1.45)^2(0.05)^2 + (2.45)^2(0.05)^2]^{1/2} = 0.15$ mag. R2 is derived by combining the partially correlated quantities (f) and (g) in quadrature. The individual random and systematic errors are then combined in quadrature to yield the total random (R) and systematic (S) errors.

Table 9. Photometry for WF2 Secondary Standards

X ^a	Y	DoPHOT		ALLFRAME	
		V	I	V	I
140.96	349.65	25.68 (0.02)	24.52 (0.02)	25.78 (0.02)	24.60 (0.02)
142.71	287.06	25.95 (0.03)	24.52 (0.02)	26.00 (0.02)	24.54 (0.01)
147.70	473.61	25.71 (0.04)	25.47 (0.02)	25.78 (0.02)	25.46 (0.03)
166.62	183.35	25.09 (0.02)	23.39 (0.02)	25.19 (0.01)	23.45 (0.01)
173.68	539.46	24.59 (0.02)	24.65 (0.04)	24.64 (0.01)	24.68 (0.01)
187.15	212.91	25.06 (0.02)	23.37 (0.02)	25.15 (0.01)	23.45 (0.01)
188.54	188.93	23.20 (0.01)	23.11 (0.02)	23.25 (0.00)	23.13 (0.01)
192.37	376.85	25.15 (0.02)	25.00 (0.02)	25.21 (0.01)	25.03 (0.02)
198.22	216.76	25.48 (0.03)	25.32 (0.04)	25.57 (0.01)	25.41 (0.03)
202.76	726.39	25.48 (0.02)	25.32 (0.05)	25.49 (0.01)	25.25 (0.04)
215.92	86.67	26.71 (0.03)	25.47 (0.03)	26.91 (0.03)	25.58 (0.03)
216.20	631.36	25.99 (0.03)	24.50 (0.02)	26.08 (0.03)	24.55 (0.02)
223.37	141.22	26.11 (0.04)	24.76 (0.03)	26.24 (0.03)	24.80 (0.02)
251.83	116.54	25.70 (0.03)	25.47 (0.04)	25.81 (0.02)	25.43 (0.03)
255.17	88.57	25.86 (0.02)	24.32 (0.02)	26.01 (0.02)	24.40 (0.02)
256.01	213.12	25.75 (0.04)	24.15 (0.03)	25.84 (0.02)	24.21 (0.01)
266.06	411.81	23.19 (0.02)	21.63 (0.01)	23.30 (0.01)	21.71 (0.00)
275.13	138.64	25.31 (0.04)	25.29 (0.05)	25.45 (0.01)	25.36 (0.03)
280.21	393.94	25.30 (0.01)	25.07 (0.04)	25.41 (0.01)	25.18 (0.03)
281.36	581.86	25.24 (0.03)	23.58 (0.02)	25.28 (0.01)	23.62 (0.01)
294.12	363.87	24.19 (0.03)	23.74 (0.01)	24.28 (0.01)	23.80 (0.01)
300.39	606.80	25.56 (0.03)	25.16 (0.03)	25.55 (0.02)	25.20 (0.03)
321.95	107.89	25.72 (0.03)	24.20 (0.04)	25.84 (0.02)	24.26 (0.01)
345.29	653.94	24.04 (0.04)	22.27 (0.02)	24.01 (0.01)	22.31 (0.01)
359.01	533.60	25.36 (0.02)	23.68 (0.04)	25.50 (0.01)	23.81 (0.01)
369.20	475.85	26.08 (0.04)	24.73 (0.03)	26.18 (0.03)	24.82 (0.02)
377.72	503.14	25.95 (0.03)	24.57 (0.03)	26.04 (0.02)	24.62 (0.01)
392.64	523.50	24.48 (0.02)	24.50 (0.02)	24.58 (0.01)	24.51 (0.02)
430.09	473.95	25.53 (0.03)	25.41 (0.03)	25.62 (0.02)	25.45 (0.03)
433.98	449.86	26.40 (0.03)	24.97 (0.03)	26.47 (0.03)	25.00 (0.02)
443.70	187.70	26.05 (0.04)	25.44 (0.03)	26.12 (0.02)	25.44 (0.04)
444.08	265.85	25.64 (0.02)	25.58 (0.04)	25.79 (0.01)	25.68 (0.03)
450.07	391.52	26.07 (0.03)	24.72 (0.04)	26.19 (0.02)	24.77 (0.03)
458.88	462.63	25.70 (0.02)	24.40 (0.03)	25.82 (0.01)	24.45 (0.01)
466.57	460.78	24.95 (0.02)	24.19 (0.03)	25.02 (0.01)	24.23 (0.01)

Table 9—Continued

X ^a	Y	DoPHOT		ALLFRAME	
		V	I	V	I
468.80	149.52	25.03 (0.02)	25.10 (0.02)	25.14 (0.01)	25.12 (0.02)
473.49	485.00	25.60 (0.02)	25.25 (0.04)	25.70 (0.01)	25.34 (0.03)
475.24	530.27	25.37 (0.02)	25.38 (0.04)	25.47 (0.02)	25.41 (0.03)
477.42	557.90	25.31 (0.02)	25.38 (0.04)	25.38 (0.01)	25.43 (0.05)
486.58	480.36	25.28 (0.02)	25.12 (0.04)	25.41 (0.01)	25.16 (0.03)
492.53	418.92	25.62 (0.02)	24.34 (0.04)	25.73 (0.02)	24.36 (0.01)
494.50	529.67	26.00 (0.02)	24.56 (0.04)	26.07 (0.02)	24.59 (0.02)
496.36	300.87	26.13 (0.03)	24.43 (0.02)	26.19 (0.02)	24.52 (0.02)
500.92	755.29	25.06 (0.03)	25.17 (0.02)	25.21 (0.01)	25.19 (0.03)
503.99	729.88	26.53 (0.04)	25.21 (0.03)	26.59 (0.03)	25.22 (0.03)
510.40	187.05	23.94 (0.02)	23.55 (0.02)	24.03 (0.01)	23.60 (0.01)
512.55	515.64	24.86 (0.02)	23.29 (0.02)	24.94 (0.01)	23.31 (0.01)
513.68	463.73	24.67 (0.01)	24.76 (0.02)	24.75 (0.01)	24.80 (0.02)
515.61	488.79	24.93 (0.03)	23.10 (0.02)	24.95 (0.01)	23.13 (0.01)
566.02	528.23	25.53 (0.03)	25.59 (0.04)	25.64 (0.02)	25.64 (0.03)
580.93	511.01	25.45 (0.02)	25.43 (0.03)	25.55 (0.01)	25.43 (0.03)
585.69	407.31	26.64 (0.05)	25.18 (0.03)	26.75 (0.04)	25.22 (0.03)

^aX and Y coordinates are derived from UT 1997 August 06 epoch.

Table 10. Photometry for WF3 Secondary Standards

X ^a	Y	DoPHOT		ALLFRAME	
		V	I	V	I
113.30	467.79	26.23 (0.04)	24.67 (0.02)	26.36 (0.03)	24.77 (0.03)
123.87	316.10	25.74 (0.02)	25.39 (0.04)	25.80 (0.02)	25.51 (0.04)
167.03	196.51	25.72 (0.02)	25.27 (0.03)	25.84 (0.03)	25.42 (0.03)
217.75	132.69	25.42 (0.02)	24.73 (0.03)	25.47 (0.02)	24.86 (0.02)
219.94	205.60	25.28 (0.02)	23.63 (0.01)	25.36 (0.01)	23.75 (0.01)
220.51	330.99	25.81 (0.05)	25.76 (0.04)	25.87 (0.03)	25.82 (0.04)
232.10	305.34	24.90 (0.03)	24.16 (0.03)	25.00 (0.02)	24.25 (0.02)
239.53	352.16	26.52 (0.04)	24.91 (0.02)	26.57 (0.03)	25.01 (0.03)
245.59	86.09	26.54 (0.04)	24.84 (0.04)	26.59 (0.03)	25.02 (0.02)
252.90	125.29	25.97 (0.02)	24.30 (0.03)	26.07 (0.02)	24.45 (0.02)
253.99	541.31	25.41 (0.01)	24.54 (0.03)	25.47 (0.01)	24.66 (0.03)
268.01	76.50	25.75 (0.04)	25.29 (0.03)	25.87 (0.02)	25.35 (0.03)
275.36	689.03	25.94 (0.03)	24.97 (0.03)	26.03 (0.02)	25.00 (0.03)
299.01	289.90	26.16 (0.03)	24.60 (0.02)	26.25 (0.02)	24.71 (0.02)
299.90	540.05	25.63 (0.04)	24.85 (0.02)	25.70 (0.02)	24.98 (0.02)
305.93	663.39	26.01 (0.03)	25.55 (0.04)	26.08 (0.02)	25.54 (0.03)
307.95	568.52	25.15 (0.02)	25.35 (0.04)	25.25 (0.01)	25.42 (0.04)
310.54	290.27	25.23 (0.03)	24.86 (0.03)	25.33 (0.01)	25.03 (0.03)
314.11	394.59	25.16 (0.03)	23.12 (0.04)	25.27 (0.02)	23.25 (0.01)
321.62	568.46	26.53 (0.05)	25.18 (0.03)	26.58 (0.03)	25.24 (0.03)
326.16	621.55	26.30 (0.03)	24.96 (0.04)	26.40 (0.03)	25.11 (0.02)
335.34	149.15	25.29 (0.03)	23.31 (0.02)	25.33 (0.01)	23.44 (0.01)
341.61	574.70	23.68 (0.02)	22.31 (0.01)	23.75 (0.01)	22.33 (0.01)
365.50	307.88	25.16 (0.02)	25.13 (0.03)	25.27 (0.01)	25.30 (0.03)
403.32	99.32	24.55 (0.04)	24.40 (0.02)	24.53 (0.01)	24.47 (0.02)
405.66	782.52	26.14 (0.04)	24.23 (0.02)	26.23 (0.03)	24.25 (0.02)
414.42	628.68	23.53 (0.02)	23.43 (0.02)	23.64 (0.01)	23.47 (0.01)
417.58	78.04	24.77 (0.03)	23.39 (0.01)	24.80 (0.01)	23.50 (0.01)
430.79	374.02	25.37 (0.02)	24.16 (0.02)	25.50 (0.02)	24.32 (0.01)
442.67	502.92	24.77 (0.02)	24.78 (0.02)	24.87 (0.01)	24.88 (0.02)
444.13	289.70	26.19 (0.03)	24.65 (0.03)	26.30 (0.02)	24.79 (0.02)
459.20	343.29	24.75 (0.02)	24.75 (0.04)	24.87 (0.01)	24.88 (0.05)
471.33	526.16	25.90 (0.02)	24.67 (0.03)	26.02 (0.02)	24.83 (0.02)
485.86	564.42	25.44 (0.04)	25.26 (0.04)	25.51 (0.01)	25.33 (0.02)
530.95	308.74	26.21 (0.03)	24.66 (0.04)	26.28 (0.03)	24.80 (0.03)

Table 10—Continued

X ^a	Y	DoPHOT		ALLFRAME	
		V	I	V	I
556.95	563.20	25.70 (0.02)	24.07 (0.02)	25.78 (0.01)	24.14 (0.01)
564.25	420.70	22.63 (0.02)	22.37 (0.01)	22.75 (0.01)	22.46 (0.01)
577.73	429.41	25.86 (0.02)	24.35 (0.02)	25.93 (0.02)	24.46 (0.02)
586.57	378.54	24.02 (0.04)	23.40 (0.02)	24.06 (0.01)	23.50 (0.01)
587.83	700.23	26.47 (0.04)	24.77 (0.03)	26.52 (0.03)	24.85 (0.03)
603.83	778.79	26.66 (0.02)	25.06 (0.05)	26.79 (0.03)	25.13 (0.04)
611.57	508.83	23.57 (0.01)	23.48 (0.01)	23.68 (0.01)	23.57 (0.01)
617.26	429.29	25.12 (0.02)	23.33 (0.03)	25.24 (0.01)	23.38 (0.01)
648.66	763.49	25.36 (0.04)	25.28 (0.03)	25.39 (0.01)	25.27 (0.03)
678.93	543.15	25.76 (0.04)	24.02 (0.02)	25.87 (0.02)	24.06 (0.01)
679.63	301.03	24.95 (0.02)	24.74 (0.04)	25.05 (0.01)	24.92 (0.03)
683.87	686.51	24.99 (0.01)	25.18 (0.04)	25.06 (0.01)	25.19 (0.04)
685.51	638.24	23.12 (0.02)	22.39 (0.03)	23.16 (0.01)	22.35 (0.01)
702.25	598.77	25.02 (0.02)	23.78 (0.03)	25.10 (0.01)	23.80 (0.01)
714.37	598.01	25.77 (0.03)	24.26 (0.02)	25.89 (0.03)	24.30 (0.01)
720.61	648.82	25.84 (0.03)	24.27 (0.04)	25.90 (0.02)	24.32 (0.02)
722.70	559.34	24.57 (0.02)	24.79 (0.04)	24.59 (0.01)	24.78 (0.02)
723.63	734.81	26.12 (0.03)	24.86 (0.04)	26.17 (0.02)	24.90 (0.02)
738.68	539.00	26.03 (0.04)	24.83 (0.03)	26.17 (0.03)	24.89 (0.03)
747.90	478.86	25.82 (0.02)	24.18 (0.03)	25.93 (0.02)	24.25 (0.02)
750.23	667.98	24.75 (0.02)	23.31 (0.03)	24.79 (0.01)	23.29 (0.01)
762.19	626.60	25.94 (0.04)	24.75 (0.02)	26.06 (0.03)	24.77 (0.02)
779.31	385.82	25.39 (0.02)	25.30 (0.04)	25.50 (0.02)	25.38 (0.03)

^aX and Y coordinates are derived from UT 1997 August 06 epoch.

Table 11. Photometry for WF4 Secondary Standards

X ^a	Y	DoPHOT		ALLFRAME	
		V	I	V	I
162.83	185.95	26.14 (0.03)	24.58 (0.03)	26.12 (0.03)	24.53 (0.01)
205.49	358.85	25.23 (0.04)	25.31 (0.04)	25.20 (0.01)	25.32 (0.02)
215.67	310.20	26.46 (0.04)	24.77 (0.03)	26.41 (0.03)	24.77 (0.02)
248.13	318.95	26.36 (0.04)	24.63 (0.03)	26.29 (0.02)	24.56 (0.01)
259.18	145.25	26.30 (0.03)	25.82 (0.04)	26.32 (0.02)	25.80 (0.04)
266.79	179.39	25.21 (0.02)	24.68 (0.04)	25.24 (0.01)	24.67 (0.02)
303.49	190.27	26.21 (0.03)	24.94 (0.03)	26.17 (0.02)	24.93 (0.03)
314.26	247.39	25.79 (0.02)	25.18 (0.04)	25.82 (0.02)	25.16 (0.03)
368.66	254.28	26.21 (0.02)	24.89 (0.03)	26.24 (0.02)	24.85 (0.02)
374.97	355.20	25.19 (0.02)	23.66 (0.03)	25.14 (0.01)	23.63 (0.01)
387.49	320.09	25.14 (0.04)	24.97 (0.04)	25.11 (0.01)	24.97 (0.02)
441.78	120.31	25.68 (0.03)	25.41 (0.04)	25.74 (0.01)	25.47 (0.05)
447.70	236.39	24.28 (0.03)	24.09 (0.05)	24.27 (0.01)	24.15 (0.01)
456.73	348.82	26.03 (0.02)	24.58 (0.04)	26.07 (0.02)	24.63 (0.02)
507.16	118.35	24.86 (0.02)	22.94 (0.04)	24.90 (0.01)	23.00 (0.01)
507.78	229.01	26.66 (0.05)	25.37 (0.04)	26.66 (0.03)	25.45 (0.04)
533.63	250.54	25.47 (0.02)	25.51 (0.05)	25.52 (0.01)	25.53 (0.04)
608.07	109.54	24.81 (0.04)	24.61 (0.02)	24.80 (0.01)	24.67 (0.02)
623.71	129.27	26.30 (0.05)	24.85 (0.03)	26.34 (0.03)	24.90 (0.02)
631.06	408.56	26.21 (0.02)	24.47 (0.04)	26.13 (0.03)	24.44 (0.01)

^aX and Y coordinates are derived from UT 1997 August 06 epoch.

Table 12. Astrometry for Other NGC 1326-A Variables

ID ^a	X ^b	Y ^b	RA(J2000) ^c	DEC(J2000) ^c	Chip
V1	454.36	248.19	3:25:08.10	−36:21:59.0	WF3
V2	667.16	544.71	3:25:11.02	−36:22:07.7	WF3
V3	342.21	325.48	3:25:07.81	−36:22:12.1	WF3
V4	83.12	482.06	3:25:07.06	−36:21:38.9	WF2
V5	662.70	743.72	3:25:12.05	−36:22:23.1	WF3
V6	172.49	609.23	3:25:07.38	−36:21:23.9	WF2
V7	382.57	126.42	3:25:07.00	−36:21:54.4	WF3
V8	743.64	632.98	3:25:11.97	−36:22:09.5	WF3
V9	274.84	193.41	3:25:06.68	−36:22:06.4	WF3
V10	768.85	334.02	3:25:10.54	−36:21:45.2	WF3
V11	103.38	389.29	3:25:03.40	−36:22:00.9	PC1
V12	105.46	99.80	3:25:04.37	−36:22:18.8	WF4
V13	680.72	136.35	3:25:08.93	−36:21:35.9	WF3
V14	312.74	475.72	3:25:05.79	−36:21:22.0	WF2
V15	365.40	218.52	3:25:07.39	−36:22:02.5	WF3
V16	280.02	329.34	3:25:05.05	−36:21:33.9	WF2
V17	318.11	725.72	3:25:09.79	−36:22:44.0	WF3

^aVariables are ordered by increasing period.

^bX and Y coordinates are derived from UT 1997 August 06 epoch.

^cRA and DEC are derived using the IRAF task `stdas.toolbox.imgtools.xy2rd`, STSDAS V2.0.1.

Notes on individual variables. V5: possible nonstellar object? V7: possible cepheid? V8: possible cepheid? V9: possible cepheid? This variable was not recovered by ALLFRAME. V11: This variable from PC chip was only measured by DoPHOT, as ALLFRAME reductions of PC1 were not performed due to the few variables found by the DoPHOT reduction and analysis. V12: possible cepheid? Variable lies near pyramid corner of chip. V13: possible cepheid? There is a strong diffraction spike in I-band due to a nearby brighter star which may contaminate F814W mag. measurement for this variable. V14: There is also a strong alias at 7.3d. V15: Only V-band photometry was obtained by DoPHOT, the star not being recovered in DoPHOT I-band photometry. V16: Noticeable aliases at 9.2d and 6.1d. V17: possible cepheid? Only V-band photometry was obtained by DoPHOT and this variable was not recovered by ALLFRAME.

Table 13. DoPHOT Mean Photometry for Other NGC 1326-A Variables

ID	P	P _{err}	χ^2_r	log P	V_{ph}	I_{ph}	V- I_{ph}
V1	>60	...	7.3	>1.78	24.65	22.73	1.92
V2	>60	...	9.0	>1.78	26.09	24.18	1.91
V3	55	(4)	12.9	1.74	24.22	23.35	0.87
V4	52.	(5)	7.5	1.72	22.87	21.44	1.43
V5	>50	...	17.8	1.70	23.38	22.77	0.61
V6	>40	...	11.9	1.60	24.80	23.06	1.74
V7	36.0	(5)	12.4	1.56	25.96	24.77	1.19
V8	30.5	(4)	4.3	1.48	26.15	25.39	0.76
V9	25.	(2)	5.4	1.40	25.89	24.86	1.03
V10	22.8	(1.5)	8.2	1.36	25.92	25.38	0.54
V11	22.8	(1.2)	8.9	1.36	26.67	25.74	0.93
V12	20.7	(2)	4.3	1.32	25.98	24.90	1.08
V13	17.4	(1)	4.5	1.24	26.34	25.23	1.11
V14	16.2	(1.5)	2.3	1.21	25.92	25.18	0.74
V15	16.0	(1)	4.5	1.20	26.79	...	...
V16	15.5	(0.5)	8.5	1.19	26.81	26.52	0.29
V17	12.6	(1)	2.9	1.10	27.09	...	...

Table 1. DoPHOT F555W Photometry at Individual Epochs

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	C1	C2	C3	C4	C5	C6	C7	C8
667.0953	25.11 0.06	25.78 0.07	26.28 0.14	25.15 0.05	25.64 0.10	26.25 0.09	26.27 0.11	26.17 0.12
670.1196	25.17 0.06	25.40 0.04	26.25 0.30	25.11 0.05	25.85 0.09	26.31 0.11	26.31 0.10	25.93 0.14
673.0080	25.27 0.07	25.29 0.06	25.16 0.08	25.17 0.05	26.00 0.08	25.92 0.07	26.31 0.10	26.30 0.14
675.9063	25.31 0.06	25.36 0.05	25.03 0.06	25.38 0.05	26.05 0.23	25.13 0.04	26.46 0.12	26.43 0.17
679.2000	25.42 0.05	25.48 0.04	25.20 0.05	25.38 0.05	26.27 0.08		25.49 0.04	26.06 0.09
684.3709	25.46 0.06		25.45 0.09	25.07 0.06	26.44 0.13	25.60 0.07	25.68 0.07	25.46 0.07
687.7434	25.65 0.06	25.62 0.06	25.64 0.06	24.86 0.05	26.39 0.10	25.60 0.05	25.72 0.07	25.56 0.08
693.3174	25.13 0.05	25.87 0.06	25.94 0.13	24.84 0.04	26.29 0.12	26.35 0.17	26.08 0.08	26.00 0.10
697.2925	24.58 0.06		26.15 0.13	24.96 0.05	25.62 0.07	26.05 0.08	26.37 0.10	26.12 0.10
704.2745	24.80 0.04	26.29 0.08	26.14 0.11	25.17 0.05	25.93 0.15	25.91 0.09	26.33 0.11	26.45 0.21
709.3821	24.83 0.04	25.44 0.07	24.82 0.07	25.25 0.06	26.00 0.10	25.27 0.05	25.47 0.08	25.19 0.06
716.1700	25.07 0.07	25.44 0.06		25.17 0.05	26.12 0.13	25.53 0.06	25.67 0.09	25.73 0.10
2450000.+	C9	C10	C11	C12	C13	C14	C15	C16
667.0953	25.85 0.08	26.13 0.11	26.41 0.15	27.01 0.22	26.34 0.13	26.24 0.11	26.49 0.14	27.01 0.18
670.1196	25.92 0.09	26.92 0.24	26.50 0.19	26.60 0.15	26.57 0.15	26.75 0.22	26.07 0.12	26.54 0.13
673.0080	26.11 0.09	26.42 0.18		26.31 0.11	26.82 0.15	27.06 0.24	26.18 0.10	27.02 0.17
675.9063	26.38 0.11	26.53 0.19	26.19 0.11	25.92 0.07	26.93 0.20	27.38 0.29	26.41 0.12	26.92 0.16
679.2000	26.44 0.09	26.81 0.19	26.27 0.12	26.21 0.09	27.00 0.17	26.83 0.14	26.88 0.15	27.47 0.24
684.3709	26.70 0.16	25.82 0.10	26.79 0.22	26.79 0.16	26.22 0.12	26.09 0.13	26.39 0.12	26.89 0.17
687.7434	25.57 0.06	26.19 0.11	26.52 0.12	26.80 0.14	26.36 0.11		25.93 0.08	27.37 0.23
693.3174	25.71 0.07	26.70 0.21	25.96 0.10	26.11 0.09	26.88 0.17	27.26 0.28	26.43 0.10	27.60 0.36
697.2925	26.04 0.09	26.63 0.16	26.20 0.14	26.27 0.09	26.71 0.47	26.85 0.17	26.66 0.13	26.79 0.14
704.2745	26.36 0.12	25.63 0.10	26.53 0.20	26.74 0.15	26.14 0.12		25.80 0.09	27.25 0.20
709.3821	26.67 0.18	25.85 0.09	26.54 0.18	27.00 0.16	26.47 0.13	27.05 0.25	25.98 0.11	27.91 0.41
716.1700	25.79 0.07	26.49 0.18	26.01 0.10	26.11 0.09	26.77 0.16	27.35 0.35	26.75 0.19	26.77 0.16
2450000.+	C17	C18	C19	V1	V2	V3	V4	V5
667.0953		26.98 0.16	27.13 0.21	24.50 0.05	26.59 0.17	24.26 0.04	23.04 0.04	23.34 0.04
670.1196	26.56 0.16	26.49 0.12	26.46 0.14	24.61 0.07	26.53 0.15	24.30 0.04		23.28 0.04
673.0080	27.30 0.25	26.33 0.10	27.07 0.20	24.47 0.05	26.51 0.14	24.37 0.03	22.96 0.04	23.32 0.04
675.9063	26.99 0.18	26.85 0.14	26.95 0.19	24.53 0.05	26.46 0.13	24.40 0.04	22.85 0.03	
679.2000	26.57 0.12	27.04 0.16	26.49 0.10	24.48 0.04	26.24 0.08	24.45 0.03	22.75 0.03	23.35 0.04
684.3709	27.10 0.24	26.55 0.13	27.03 0.17	24.50 0.05	26.30 0.09	24.46 0.03	22.74 0.03	23.32 0.03
687.7434	27.44 0.30	27.25 0.21	26.36 0.10	24.59 0.05	26.10 0.10	24.40 0.06	22.72 0.03	23.41 0.03
693.3174	26.17 0.09	26.52 0.12	27.36 0.23	24.61 0.05	25.98 0.07	24.42 0.04	22.75 0.03	23.34 0.03
697.2925	27.04 0.20	26.57 0.11	26.52 0.12	24.66 0.04	25.90 0.08	24.35 0.03	22.80 0.03	23.47 0.03
704.2745	26.11 0.12	26.39 0.12		24.70 0.05	25.86 0.08	24.02 0.03	22.98 0.04	23.54 0.03
709.3821	26.58 0.15	26.84 0.14	26.41 0.12	24.76 0.06	25.74 0.08	23.97 0.04	23.02 0.04	23.60 0.03
716.1700	26.29 0.10	26.55 0.14	26.89 0.16	24.95 0.05	25.87 0.10	24.01 0.04	23.07 0.04	
2450000.+	V6	V7	V8	V9	V10	V11	V12	V13
667.0953	24.86 0.06	26.06 0.12	26.47 0.14	26.24 0.12	25.59 0.08	27.00 0.17		26.76 0.22
670.1196	25.00 0.08	26.24 0.16		26.18 0.14	26.07 0.11	27.18 0.22	25.98 0.11	27.03 0.30
673.0080	24.96 0.06	26.23 0.15	26.16 0.10	25.70 0.06	26.17 0.10	26.54 0.14	26.00 0.10	26.75 0.21
675.9063	24.90 0.04	26.42 0.20	26.08 0.12	25.69 0.08	26.25 0.13	26.27 0.10	26.32 0.12	26.05 0.13
679.2000	24.84 0.04	26.39 0.14	25.98 0.08	25.84 0.07	26.17 0.09	26.43 0.11	26.20 0.11	26.17 0.11
684.3709	24.87 0.05	25.64 0.12	25.85 0.10	25.78 0.09	25.78 0.12	26.60 0.12	25.71 0.07	26.66 0.25
687.7434	24.75 0.04	25.59 0.09	26.11 0.09	26.01 0.08	25.54 0.08	26.91 0.10	25.87 0.08	26.67 0.20
693.3174	24.78 0.05	26.00 0.13	26.48 0.13	26.12 0.10	26.08 0.11	27.38 0.20	26.03 0.10	26.00 0.13
697.2925	24.76 0.04	25.88 0.11		26.01 0.09		26.27 0.10	26.23 0.11	26.12 0.14
704.2745	24.79 0.06	26.00 0.12	26.20 0.13	25.84 0.10	26.04 0.12	26.91 0.15	25.85 0.09	26.55 0.21
709.3821	24.65 0.05	26.44 0.16	25.95 0.10		25.75 0.09	27.05 0.20	25.88 0.11	26.06 0.13
716.1700			26.07 0.11	26.21 0.12	26.04 0.12	27.24 0.21	26.31 0.17	26.23 0.14

Table 1—Continued

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	V14	V15	V16	V17				
667.0953	25.93 0.10	26.38 0.15						
670.1196	25.87 0.11	26.75 0.33	27.02 0.20	27.56 0.34				
673.0080	25.95 0.09	27.99 0.21	26.95 0.18	27.07 0.18				
675.9063		27.55 0.43	27.30 0.22	26.81 0.16				
679.2000	26.03 0.08	27.13 0.25	26.41 0.11	26.89 0.15				
684.3709	25.70 0.06	26.36 0.16						
687.7434		26.69 0.20		26.95 0.17				
693.3174	25.87 0.08	27.25 0.30	27.24 0.20	27.34 0.25				
697.2925	25.97 0.08	26.38 0.14	26.46 0.10	27.12 0.16				
704.2745	25.99 0.10	27.11 0.28	27.12 0.20	27.17 0.24				
709.3821	26.15 0.11	27.32 0.34	26.80 0.14	27.47 0.32				
716.1700	26.03 0.10	26.34 0.17	26.79 0.15	26.99 0.29				

Table 2. DoPHOT F814W Photometry at Individual Epochs

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	C1	C2	C3	C4	C5	C6	C7	C8
667.1765	24.20 0.05	24.94 0.07	25.19 0.10	24.09 0.05	25.20 0.11	25.13 0.07	25.07 0.09	24.87 0.10
670.2009	24.10 0.05	24.76 0.07	25.17 0.09	24.26 0.05	24.93 0.08	25.36 0.09	25.44 0.12	25.20 0.12
673.0917	24.16 0.05	24.66 0.05	24.67 0.10	24.20 0.05	24.99 0.08	25.15 0.08	25.27 0.08	
684.4542	24.54 0.05	24.69 0.06	24.68 0.07	24.14 0.05	25.00 0.09	24.76 0.08	24.93 0.06	24.93 0.12
693.4007	24.29 0.05	24.82 0.05	25.08 0.13	24.04 0.06	25.32 0.10	25.00 0.08	25.09 0.08	25.14 0.12
704.3557	24.06 0.05	25.15 0.08	25.34 0.13	24.29 0.06	24.91 0.09	25.02 0.07	25.46 0.13	25.29 0.13
709.4633	24.03 0.05	24.81 0.07	24.47 0.07	24.38 0.07	24.95 0.08	24.62 0.06	24.83 0.08	24.58 0.06
716.2515	24.14 0.05	24.78 0.07	24.44 0.07	24.37 0.06	25.48 0.13	24.75 0.06	26.02 0.08	24.96 0.11
2450000.+	C9	C10	C11	C12	C13	C14	C15	C16
667.1765	25.31 0.09	25.32 0.20	25.04 0.11	25.90 0.13	25.32 0.15	25.40 0.16	25.55 0.12	25.98 0.17
670.2009	25.13 0.09	25.24 0.14	24.97 0.12	25.98 0.15	25.41 0.12	25.68 0.19	25.25 0.11	26.16 0.20
673.0917	25.26 0.09	25.42 0.15	24.99 0.11	25.99 0.15	25.26 0.11	25.81 0.19	25.13 0.09	26.20 0.19
684.4542	25.67 0.14	25.05 0.12	25.20 0.12	25.85 0.17	25.52 0.16	25.46 0.14	25.51 0.12	26.42 0.22
693.4007	25.04 0.07	25.56 0.15	25.14 0.10	25.53 0.10	25.30 0.11	26.39 0.30	25.36 0.09	
704.3557	25.65 0.14	25.09 0.11	25.17 0.13	25.80 0.14	25.20 0.11		25.06 0.09	26.08 0.16
709.4633	25.65 0.13	24.91 0.15	24.85 0.10	26.56 0.25	25.36 0.12		25.08 0.09	
716.2515	25.13 0.09	25.44 0.17	24.71 0.11	25.60 0.11	25.56 0.17	26.31 0.31	25.64 0.16	26.50 0.26
2450000.+	C17	C18	C19	V1	V2	V3	V4	V5
667.1765	25.85 0.20	26.30 0.23	26.16 0.21	22.76 0.05	24.53 0.07	23.37 0.05	21.53 0.05	22.58 0.05
670.2009	26.30 0.29	26.03 0.17	26.12 0.20	22.73 0.05	24.62 0.07	23.26 0.05	21.46 0.05	22.82 0.05
673.0917	25.92 0.21	25.56 0.15	26.48 0.28	22.67 0.05	24.59 0.07	23.40 0.05	21.47 0.05	
684.4542	26.27 0.27	25.68 0.13	26.25 0.22	22.73 0.05	24.39 0.05	23.52 0.05	21.48 0.05	22.87 0.06
693.4007	25.55 0.17	25.75 0.13	26.69 0.26	22.75 0.05	24.12 0.08	23.54 0.05	21.43 0.05	
704.3557	25.84 0.21	26.35 0.24	26.64 0.26	22.82 0.05	24.12 0.06	23.34 0.05	21.52 0.05	22.75 0.05
709.4633	25.84 0.17	26.18 0.24	25.96 0.15	22.82 0.05	24.15 0.06	23.23 0.05	21.43 0.05	22.72 0.05
716.2515	26.16 0.27	25.90 0.14	26.80 0.29	22.74 0.05	24.04 0.05	23.25 0.05	21.40 0.05	23.05 0.05
2450000.+	V6	V7	V8	V9	V10	V11	V12	V13
667.1765	23.19 0.05	24.97 0.09	25.77 0.17	25.05 0.09	25.06 0.15		24.94 0.10	25.53 0.19
670.2009	23.08 0.05	25.13 0.13	25.59 0.15	25.22 0.13	25.45 0.16	26.05 0.19	24.89 0.08	25.85 0.28
673.0917	23.30 0.05	25.23 0.17	25.19 0.11	24.72 0.07	25.61 0.18	25.58 0.11	25.06 0.11	25.30 0.15
684.4542	23.17 0.05	25.01 0.11	25.26 0.12	24.79 0.08	25.11 0.14	25.75 0.16	24.75 0.08	26.02 0.13
693.4007	23.09 0.05	24.51 0.11	25.30 0.13	25.36 0.11	25.54 0.18	25.94 0.22	25.18 0.10	25.05 0.12
704.3557	23.05 0.08	24.88 0.11	25.25 0.11	24.79 0.09	25.60 0.17	25.67 0.19	24.75 0.08	25.20 0.15
709.4633	22.96 0.05	25.30 0.14	25.32 0.14	24.65 0.08	25.44 0.18	25.94 0.17	24.97 0.09	
716.2515	23.11 0.05	25.39 0.13	25.61 0.15	25.15 0.12	25.67 0.12	26.09 0.18	25.17 0.10	25.51 0.17
2450000.+	V14	V15	V16	V17				
667.1765	25.16 0.08							
670.2009	25.08 0.09		26.19 0.20					
673.0917	25.19 0.08		26.87 0.46					
684.4542	25.10 0.08		26.87 0.36					
693.4007	25.36 0.10		26.98 0.44					
704.3557	25.16 0.10		26.52 0.27					
709.4633	25.43 0.11		26.20 0.20					
716.2515	25.16 0.09		26.27 0.15					

Table 16. ALLFRAME F555W Photometry at Individual Epochs

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	C1	C2	C3	C4	C5	C6	C7	C8
667.0953	25.17 0.04	25.84 0.05	26.22 0.11	25.19 0.04	25.30 0.14	26.22 0.08	26.34 0.07	
670.1196	25.25 0.03	25.52 0.04	26.18 0.08	25.19 0.04	25.94 0.06	26.38 0.08	26.22 0.13	
673.0080	25.37 0.04	25.39 0.04	25.02 0.03	25.26 0.06	26.02 0.08	25.88 0.05	26.36 0.11	
675.9063	25.41 0.03	25.50 0.04	25.04 0.04	25.41 0.04	26.10 0.10	25.20 0.03	26.53 0.06	
679.2000	25.51 0.08	25.54 0.03	25.19 0.04	25.42 0.03	26.30 0.07	25.52 0.09	25.57 0.04	
684.3709	25.61 0.05	25.68 0.05	25.40 0.05	25.11 0.03	26.58 0.08	25.62 0.04	25.77 0.05	
687.7434	25.73 0.05	25.77 0.08	25.60 0.07	24.90 0.03	26.49 0.08	25.71 0.04	25.77 0.04	
693.3174	25.21 0.04	25.97 0.07	25.75 0.13	24.93 0.07	26.54 0.11	26.10 0.07	26.06 0.05	
697.2925	24.69 0.03	26.08 0.08	26.09 0.07	25.10 0.04	25.65 0.05	26.12 0.06	26.38 0.06	
704.2745	24.90 0.04	26.34 0.06	26.48 0.08	25.27 0.04	25.82 0.05	25.92 0.07	26.41 0.09	
709.3821	25.03 0.05	25.50 0.06	24.63 0.33	25.36 0.05	26.10 0.07	25.37 0.04	25.59 0.06	
716.1700	25.17 0.03	25.53 0.04	25.34 0.04	25.27 0.04	26.28 0.10	25.60 0.04	25.79 0.06	
2450000.+	C9	C10	C11	C12	C13	C14	C15	C16
667.0953	25.91 0.06		26.60 0.14	27.08 0.15	26.26 0.09	26.29 0.08	26.65 0.11	27.15 0.13
670.1196	26.03 0.06		26.35 0.13	26.79 0.11	26.42 0.10	26.63 0.14	26.16 0.08	26.72 0.09
673.0080	26.21 0.07		25.88 0.09	26.52 0.10	26.80 0.10	26.81 0.16	26.21 0.08	27.25 0.15
675.9063	26.54 0.08		26.14 0.08	26.23 0.08	26.87 0.16	27.00 0.14	26.41 0.06	27.25 0.15
679.2000	26.35 0.06		26.32 0.10	26.35 0.06	26.90 0.12	26.86 0.13	26.87 0.11	27.54 0.32
684.3709	26.76 0.09		26.88 0.18	26.88 0.10	26.33 0.08	26.29 0.10	26.54 0.09	27.22 0.17
687.7434	25.78 0.06		26.67 0.13	26.88 0.12	26.42 0.08	26.37 0.09	25.94 0.08	27.56 0.32
693.3174	25.85 0.05		25.97 0.10	26.21 0.06	27.04 0.19	27.29 0.28	26.55 0.07	27.23 0.18
697.2925	26.14 0.11		26.15 0.09	26.35 0.07	26.95 0.14	26.89 0.11	26.69 0.07	26.80 0.13
704.2745	26.47 0.10		26.35 0.15	26.77 0.09	26.25 0.08	25.95 0.09	25.96 0.07	27.71 0.33
709.3821	26.75 0.11		26.61 0.16	26.85 0.14	26.48 0.08	26.71 0.14	26.19 0.11	
716.1700	25.84 0.06		26.09 0.07	26.22 0.06	26.79 0.12	27.35 0.21	26.89 0.13	27.13 0.16
2450000.+	C17	C18	C19	V1	V2	V3	V4	V5
667.0953	26.42 0.09	26.87 0.13	27.02 0.16	24.58 0.04	26.80 0.15	24.33 0.05	22.79 0.02	23.31 0.02
670.1196	26.65 0.13	26.54 0.11	26.52 0.10	24.60 0.04	26.67 0.12	24.34 0.03	22.76 0.03	23.29 0.02
673.0080	27.16 0.15	26.37 0.08	27.04 0.12	24.52 0.03	26.61 0.08	24.46 0.03	22.76 0.03	23.27 0.03
675.9063	26.82 0.18	26.94 0.15	26.96 0.14	24.60 0.03	26.61 0.07	24.49 0.03	22.74 0.02	23.30 0.04
679.2000	26.67 0.11	26.91 0.14	26.59 0.10	24.56 0.03	26.44 0.09	24.51 0.02	22.71 0.03	23.29 0.02
684.3709	26.84 0.14	26.58 0.11	27.06 0.10	24.62 0.03	26.40 0.08	24.60 0.03	22.76 0.02	23.37 0.02
687.7434	27.04 0.31	27.19 0.15	26.43 0.07	24.64 0.03	26.34 0.09	24.59 0.03	22.72 0.03	23.31 0.04
693.3174	26.26 0.13	26.51 0.09	27.41 0.16	24.68 0.06	26.21 0.11	24.49 0.05	22.71 0.02	23.41 0.04
697.2925	26.82 0.11	26.68 0.09	26.58 0.08	24.76 0.03	26.05 0.06	24.38 0.02	22.70 0.02	23.45 0.03
704.2745	26.20 0.10	26.49 0.09	27.57 0.19	24.79 0.04	26.01 0.07	24.09 0.02	22.76 0.03	23.57 0.02
709.3821	26.75 0.11	26.86 0.10	26.58 0.16	24.77 0.04	25.88 0.06	24.06 0.02	22.73 0.04	23.70 0.04
716.1700	26.50 0.10	26.61 0.09	27.04 0.15	24.85 0.04	25.93 0.06	24.09 0.02	22.74 0.02	23.93 0.05
2450000.+	V6	V7	V8	V9	V10	V11	V12	V13
667.0953	25.05 0.06	26.18 0.10	26.52 0.10		25.68 0.08		26.05 0.07	27.19 0.20
670.1196	24.98 0.04	26.36 0.12	26.65 0.13		26.12 0.08		25.94 0.06	26.80 0.22
673.0080	25.01 0.04	26.30 0.12	26.16 0.07		26.27 0.08		26.05 0.08	26.86 0.14
675.9063	25.01 0.03	26.46 0.10	26.06 0.05		26.38 0.07		26.31 0.10	26.17 0.08
679.2000	24.91 0.03	26.46 0.10	26.03 0.06		26.28 0.07		26.20 0.06	26.41 0.07
684.3709	24.87 0.06	25.72 0.06	25.97 0.06		25.72 0.05		25.74 0.06	26.82 0.13
687.7434	24.80 0.03	25.64 0.05	26.26 0.08		25.72 0.06		25.82 0.05	27.06 0.20
693.3174	24.79 0.03	25.78 0.18	26.29 0.07		26.38 0.16		26.04 0.11	26.13 0.09
697.2925	24.70 0.07	25.91 0.06	26.31 0.08		26.63 0.11		26.26 0.07	26.33 0.10
704.2745	24.70 0.03	26.01 0.08	26.23 0.08		26.19 0.09		25.83 0.06	26.66 0.13
709.3821	24.69 0.05	26.46 0.12	25.87 0.07		25.91 0.07		25.85 0.09	26.22 0.10
716.1700	24.71 0.05	26.68 0.19	26.08 0.07		26.16 0.09		26.18 0.08	26.45 0.10

Table 16—Continued

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	V14	V15	V16	V17				
667.0953	25.77 0.06	26.40 0.10						
670.1196	25.82 0.06	26.25 0.21	27.05 0.17					
673.0080	25.91 0.06	26.67 0.19	27.13 0.13					
675.9063	26.02 0.09	27.11 0.19	27.14 0.19					
679.2000	26.07 0.06	27.05 0.33	26.51 0.09					
684.3709	25.86 0.08	26.61 0.13	27.20 0.18					
687.7434	25.94 0.06	26.67 0.11	27.08 0.15					
693.3174	26.02 0.07	27.18 0.31	27.27 0.15					
697.2925	26.04 0.06	26.37 0.10	26.58 0.08					
704.2745	25.85 0.08	27.00 0.24	27.22 0.15					
709.3821	26.03 0.07		26.72 0.16					
716.1700	25.84 0.07	26.71 0.17	27.00 0.12					

Table 17. ALLFRAME F814W Photometry at Individual Epochs

JD	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m	m $\pm$ Δ m
2450000.+	C1	C2	C3	C4	C5	C6	C7	C8
667.1765	24.19 0.05	24.91 0.06	25.18 0.06	24.09 0.04	25.00 0.11	25.32 0.08	25.08 0.08	
670.2009	24.16 0.03	24.70 0.11	25.10 0.07	24.26 0.04	25.21 0.10	25.33 0.07	25.45 0.08	
673.0917	24.22 0.07	24.66 0.03	24.30 0.12	24.18 0.04	25.04 0.06	25.10 0.05	25.30 0.05	
684.4542	24.64 0.05	24.70 0.05	24.62 0.05	24.18 0.04	25.33 0.14	24.75 0.04	24.89 0.06	
693.4007	24.35 0.03	24.83 0.03	24.86 0.10	24.09 0.03	25.41 0.08	25.09 0.07	25.01 0.05	
704.3557	24.13 0.07	25.14 0.06	25.24 0.15	24.30 0.04	24.98 0.05	24.99 0.05	25.60 0.12	
709.4633	24.08 0.04	24.78 0.05	24.50 0.06	24.36 0.05	25.07 0.08	24.68 0.05	25.01 0.07	
716.2515	24.18 0.06	24.85 0.06	24.38 0.06	24.42 0.04	25.56 0.09	24.74 0.05	25.07 0.06	
2450000.+	C9	C10	C11	C12	C13	C14	C15	C16
667.1765	25.21 0.06		25.24 0.09	25.82 0.11	25.33 0.09	25.49 0.12	25.59 0.11	26.04 0.10
670.2009	25.17 0.05		25.08 0.11	25.98 0.10	25.56 0.10	25.72 0.11	25.25 0.07	26.23 0.12
673.0917	25.27 0.07		25.03 0.08	25.98 0.15	25.42 0.08	25.95 0.14	25.19 0.09	26.34 0.12
684.4542	25.63 0.12		25.29 0.10	25.71 0.09	25.56 0.10	25.73 0.17	25.48 0.07	26.11 0.13
693.4007	25.04 0.05		25.08 0.07	25.54 0.08	25.48 0.12	26.22 0.28	25.37 0.08	26.66 0.20
704.3557	25.67 0.16		25.39 0.10	25.82 0.11	25.51 0.11	25.45 0.12	25.11 0.06	25.93 0.15
709.4633	25.63 0.14		25.09 0.09	26.41 0.20	25.44 0.10	25.61 0.12	25.19 0.10	
716.2515	25.09 0.07		25.02 0.09	25.46 0.07	25.96 0.14	26.08 0.20	25.73 0.10	26.31 0.27
2450000.+	C17	C18	C19	V1	V2	V3	V4	V5
667.1765	25.79 0.15	26.03 0.15	25.85 0.13	22.77 0.03	24.61 0.04	23.39 0.03	21.44 0.06	22.51 0.03
670.2009	26.34 0.20	25.80 0.17	25.83 0.11	22.77 0.02	24.68 0.05	23.52 0.07	21.46 0.02	22.48 0.02
673.0917	25.93 0.11	25.64 0.11	26.18 0.25	22.75 0.02	24.64 0.05	23.49 0.03	21.49 0.02	22.44 0.03
684.4542	26.13 0.16	25.64 0.11	26.39 0.14	22.82 0.02	24.42 0.04	23.60 0.03	21.48 0.02	22.43 0.05
693.4007	25.89 0.18	25.81 0.11	26.55 0.17	22.83 0.03	24.25 0.04	23.55 0.03	21.41 0.03	22.40 0.03
704.3557	25.91 0.18	25.79 0.15	26.68 0.23	22.85 0.03	24.19 0.04	23.37 0.08	21.45 0.04	22.49 0.05
709.4633	25.97 0.13	26.01 0.26	26.11 0.18	22.87 0.03	24.21 0.03	23.33 0.03	21.44 0.02	22.64 0.03
716.2515	25.92 0.17	25.99 0.11	26.08 0.21	22.91 0.04	24.18 0.05	23.34 0.03	21.43 0.03	22.71 0.03
2450000.+	V6	V7	V8	V9	V10	V11	V12	V13
667.1765	23.16 0.02	25.02 0.08	25.93 0.14		25.16 0.09		24.87 0.07	25.50 0.15
670.2009	23.18 0.03	25.18 0.10	25.54 0.07		25.50 0.11		24.95 0.07	25.85 0.19
673.0917	23.18 0.03	25.25 0.10	25.18 0.10		25.71 0.12		25.05 0.05	25.35 0.11
684.4542	23.02 0.03	25.13 0.10	25.24 0.06		25.12 0.09		24.75 0.05	25.27 0.13
693.4007	23.05 0.03	24.90 0.16	25.17 0.08		25.51 0.08		25.11 0.09	25.11 0.08
704.3557	23.02 0.03	24.99 0.11	25.16 0.09		25.57 0.12		24.81 0.05	25.40 0.12
709.4633	22.98 0.04	25.39 0.11	25.05 0.07		25.38 0.12		25.03 0.06	25.11 0.13
716.2515	22.97 0.02	25.37 0.11	25.32 0.08		25.61 0.13		25.09 0.07	25.37 0.13
2450000.+	V14	V15	V16	V17				
667.1765	25.13 0.07	26.17 0.23						
670.2009	25.12 0.06		26.16 0.14					
673.0917	25.25 0.07	27.11 0.43	26.61 0.32					
684.4542	25.15 0.05	25.88 0.22	26.60 0.21					
693.4007	25.35 0.07	26.39 0.31	26.19 0.23					
704.3557	25.31 0.11	26.56 0.36	26.56 0.29					
709.4633	25.30 0.08	26.28 0.27	25.61 0.24					
716.2515	25.15 0.08	26.11 0.22	26.28 0.11					